

Patient information factsheet

Do **not** eat or drink anything until

to normal. You can do this by sipping some cold water.

Repeat until swallowing feels normal.

- If you can feel yourself swallowing, you can eat and drink normally.

Endoscopic retrograde cholangio pancreatography (ERCP) – sedation (E1) aftercare advice

You have had a procedure called an endoscopic ret (ERCP) today:	rograde cholangio pancreatograph
This factsheet explains what to expect after your pror concerns, please contact us using the details at	
Before you leave the endoscopy unit Before you leave the endoscopy unit, we will give you: a copy of your consent form an advice sheet on what you should and should not follow-up advice a prescription (if needed)	do after your procedure
We will also explain what we have done today and wha	t we have found.
Patient sticker	Throat spray
	Sedation
	Sphincterotomy
	Stone removal
	Stent inserted
	Bile duct dilated
	Diathermy used
Table completed and advice given by: Sedation As you have had a sedative injection as part of your pro-	
capabilities may be temporarily affected. For your own should follow the advice below:	safety and the safety of others, you

If swallowing does not feel normal yet, wait a further 15 to 30 minutes and try again.

Before you eat anything, it is important that you first check your ability to swallow is back

Patient information factsheet

- When swallowing feels normal again, you may resume taking your usual medications (unless you have been told otherwise).
- Do **not** drive any type of motor vehicle or ride any type of bicycle for at least 24 hours after your procedure.
- Do **not** operate any type of electrical or mechanical equipment/machinery for at least 24 hours after your procedure.
- Do not sign any legally binding documents for at least 24 hours after your procedure.
- Do **not** drink any alcohol for at least 24 hours after your procedure.
- Rest guietly for the remainder of the day.
- You must have a responsible adult stay with you overnight.
- You should **not** be responsible for young children, disabled or dependent relatives for at least 24 hours after your procedure.

Side effects

After the procedure, it is normal to experience the following side effects:

- windy, cramp-like pain
- a sore throat

These side effects should settle within 24 to 48 hours.

If the pain does not settle or you are at all concerned about your recovery, please contact your general practitioner (GP) or our endoscopy unit using the details at the end of this factsheet.

When to seek urgent medical help

Go to your nearest emergency department if you experience any:

- unexpected or severe pain in your abdomen
- bleeding

In an emergency, call 999 for an ambulance.

Please remember to take this factsheet with you.

Follow-up care

If you need any follow-up appointments at the hospital, we will send you a letter by post.

If you need to see a GP in the two weeks after your procedure, it is important that you take this factsheet with you.

Contact us

If you have any questions or concerns while in the endoscopy unit, please speak to a member of the healthcare team caring for you.

If you have any non-urgent questions or concerns after leaving hospital, please contact your GP or our endoscopy unit for advice.

Endoscopy unit

Telephone: **023 8120 6066** (every day, 9am to 6pm)

Outside of these hours, call NHS 111 for advice.

Patient information factsheet

If you are a patient at one of our hospitals and need this document translated, or in another format such as easy read, large print, Braille or audio, please telephone **0800 484 0135** or email **patientsupporthub@uhs.nhs.uk**

For help preparing for your visit, arranging an interpreter or accessing the hospital, please visit www.uhs.nhs.uk/additionalsupport

Join our family of charity supporters with a monthly donation! It's a wonderful way to show your ongoing support of our patients and staff.





Scan the QR code or visit southamptonhospitalscharity.org/donate