

Patient information factsheet

Having a colonoscopy procedure under General Anaesthetic/Propofol sedation

Information for people with diabetes

We have written this factsheet to help you understand how to manage your diabetes before and after an endoscopic procedure such as a colonoscopy. It is important that you read and follow the instructions in this factsheet.

If you have any further questions or are unclear about any of the information in this factsheet, please contact the endoscopy unit.

Adjusting your diabetes treatments

When you go into hospital for an endoscopic procedure, you may need to adjust your diabetes treatment. This could affect your blood glucose levels because of the changes in your routine. Please do not worry about this as the levels should return to normal within 24 to 48 hours after the procedure.

If you normally check your blood glucose levels with a meter or alternative device (glucose sensor, for example) please continue to do so. You may need to test them more often. We suggest that you test before each meal, before going to bed and whenever you feel that your blood glucose levels are falling (hypoglycaemia or 'hypo').

Symptoms and signs of hypoglycaemia

These can include:

- sweating
- shaking
- blurred vision
- extreme hunger
- drowsiness
- light headedness
- slurred speech
- muddled thinking

If your blood glucose level falls below 4 mmol/L take something sugary immediately, such as:

- three to five glucose tablets
- three jelly baby sweets
- 150mls (quarter pint) smooth orange juice (no bits)

Wait 15 minutes and re-test your blood glucose levels. If they remain below 4mmol/L, repeat the initial hypo treatment.

On the day of your appointment please bring your diabetes tablets and/or insulin with you so that you can continue with it as soon as possible after your procedure.

You should have help and support from another adult while you are fasting or taking bowel preparation. They need to be able to recognise if you are having a hypo (low blood sugar) as this is more likely at this time.

Before a colonoscopy (or if having a gastroscopy at the same time as a colonoscopy)

If your diabetes is managed by diet alone

No changes are required.

If you use tablets and/or non-insulin injectable medications

Examples of non-insulin injectables: Liraglutide, Dulaglutide, Semaglutide, Tirzepatide

The day before and on the day of the procedure:

- **Stop** all **oral** diabetes medication
- **Continue** non-insulin injectable medication

After the procedure:

Once you are eating and drinking normally, you can continue your usual medications/treatment at your normal doses (**except dapagliflozin, canagliflozin, empagliflozin and ertugliflozin which should restart 48 hours after the procedure** and when eating and drinking normally).

If you use insulin for diabetes

Follow the instructions relevant to your insulin treatment:

Insulin and frequency	Two days before procedure	Day before procedure	Day of procedure
Once daily insulin (evening) - Tresiba	Take 80% of usual dose at the usual time.	Take 80% of usual dose at the usual time.	Take usual dose if eating normally in the evening.
Once daily insulin (morning) - Tresiba	Take 80% of usual dose at the usual time.	Take 80% of usual dose at the usual time.	Take 80% of usual dose - return to usual dose tomorrow.
Once daily (evening) (e.g. Lantus, Levemir, Abasaglar Insulatard or Humulin I, Toujeo)	No change	Take 80% of usual dose at the usual time.	Take 80% of your normal dose in the evening after the procedure.
Once daily (morning) as per examples above	No change	Take 80% of usual dose at the usual time.	Take 80% of your normal dose on the morning of your procedure.
Twice daily (e.g. Novomix 30, Humulin M3, Humalog Mix 25 or 50, Lantus, Levemir)	No change	Take half of usual insulin doses with a drink (containing 3 teaspoons of sugar).	Take 25% usual dose of insulin in the morning, and sugary drinks (containing 3 teaspoons of sugar).
3 to 5 injections daily (e.g. NovoRapid, Humalog, Actrapid, Humulin S, Apidra, Fiasp with long acting insulin)	No change	Do not take any rapid insulin.	Do not take any rapid insulin.

Your normal insulin dose can be resumed the day after the procedure (assuming you are able to eat and drink normally).

Patient information factsheet

If your diabetes is managed by personal insulin pump

Please inform your specialist pump team before admission.

The day before your procedure:

Continue with your usual basal rates and continue to bolus depending on carbohydrate intake. (Be mindful of what you are allowed to eat and drink for hypos and when you need to start your solid fast).

On the day of your procedure:

Continue with usual insulin infusion rate (or temporary basal rate if you have been previously trained to set this for fasting states) and continue to bolus for meals as you would do once you start eating and drinking normally.

Contact us

If you have any questions about anything related to this factsheet you can contact:

Endoscopy pre-assessment team on telephone: **07825 11807** or **07769 364198**
(Monday to Friday, 8am to 4pm).

Alternatively, you can call **the Endoscopy department direct** on telephone: **023 8120 6066**.

If for any reason your procedure is cancelled or rescheduled please contact the Endoscopy pre-assessment team on the number above.

The University Hospital Southampton NHS Foundation Trust endoscopy and adult diabetes worked together to produce this guidance.

If you are a patient at one of our hospitals and need this document translated, or in another format such as easy read, large print, Braille or audio, please telephone **0800 484 0135** or email **patientsupporthub@uhs.nhs.uk**

For help preparing for your visit, arranging an interpreter or accessing the hospital, please visit **www.uhs.nhs.uk/additionalsupport**

Join our family of charity supporters with a monthly donation!
It's a wonderful way to show your ongoing support of our patients and staff.

Scan the QR code or visit **southamptonhospitalscharity.org/donate**



**Southampton
Hospitals
Charity**

Charity Registration Number 1051543

